

AD-A031 956

WISCONSIN UNIV MADISON MATHEMATICS RESEARCH CENTER

F/G 12/1

POSITIVE EIGENFUNCTIONS FOR A CLASS OF SECOND-ORDER ELLIPTIC EQ--ETC(U)

SEP 76 P CLEMENT

DAAG29-75-C-0024

NL

UNCLASSIFIED

MRC-TSR-1673


1 OF 1  
AD  
A031956



END

DATE  
FILMED

1 77



ADA031956

MRC Technical Summary Report 1673

POSITIVE EIGENFUNCTIONS FOR A CLASS  
OF SECOND-ORDER ELLIPTIC EQUATIONS  
WITH STRONG NONLINEARITY

Philippe Clément

**Mathematics Research Center  
University of Wisconsin-Madison  
610 Walnut Street  
Madison, Wisconsin 53706**

September 1976

(Received March 1, 1976)

1473



**Approved for public release  
Distribution unlimited**

Sponsored by

U. S. Army Research Office  
P. O. Box 12211  
Research Triangle Park  
North Carolina 27709

UNIVERSITY OF WISCONSIN - MADISON  
MATHEMATICS RESEARCH CENTER

POSITIVE EIGENFUNCTIONS FOR A CLASS OF SECOND-ORDER  
ELLIPTIC EQUATIONS WITH STRONG NONLINEARITY †

Philippe Clément

Technical Summary Report #1673  
September 1976

ABSTRACT

Let  $\Omega$  be a bounded domain of  $\mathbb{R}^N$ . We consider the equation  $\mathcal{A}u(x) + f(x, u(x)) = \lambda u(x)$ ,  $\int_{\Omega} |u|^2(x) dx = R^2 > 0$ , where  $\mathcal{A}$  is a second-order quasilinear elliptic operator whose coefficients have polynomial growth and  $f$  essentially satisfies a sign condition. The existence of positive and negative solutions is proved.

AMS(MOS) Subject Classification - 35J20, 35D05, 47H15.

Key Words - Nonlinear eigenvalue problem, Strong nonlinearity, Positive solution.

Work Unit Number 1 - Applied Analysis

† Supported by the Fonds National Suisse de la Recherche Scientifique.

Sponsored by the United States Army under Contract No. DAAG29-75-C-0024.

# POSITIVE EIGENFUNCTIONS FOR A CLASS OF SECOND-ORDER ELLIPTIC EQUATIONS WITH STRONG NONLINEARITY†

Philippe Clément

The aim of this note is to prove the existence of positive and negative solutions of the eigenvalue problem;

$$(1) \quad Au = \lambda u \quad \text{with the } L^2 \text{ norm of } u, \|u\| \text{ being a prescribed constant } R > 0.$$

Here  $A$  is a second-order elliptic operator defined on a bounded domain  $\Omega$  of  $\mathbb{R}^n$ , with a strong nonlinearity in its lowest order term. We assume that  $A$  has a variational structure. The corresponding problem

(2)  $Au = f$  has been considered by several people [1], [2], [3], [4]. In these papers, only a divergence structure condition for  $A$  is assumed, however for the eigenvalue problem, such a hypothesis is too weak in general. Moreover [1], [2], [4] also deal with equations of higher order. In such situations, one can expect the existence of infinitely many distinct pairs of solutions with a prescribed norm, provided that  $A$  is odd, but not necessarily the existence of positive or negative solutions. Therefore it is convenient to consider the second-order case in itself.

## 1. Statement of the results

Let  $\Omega$  be a bounded domain of  $\mathbb{R}^N$ .

$$a: \Omega \times \mathbb{R} \times \mathbb{R}^N \rightarrow \mathbb{R} \quad \text{measurable for all } (t, \xi) \in \mathbb{R} \times \mathbb{R}^N \text{ and } C^1(\mathbb{R} \times \mathbb{R}^N; \mathbb{R})$$

for almost all  $x \in \Omega$ , satisfying the following conditions:

---

† Supported by the Fonds National Suisse de la Recherche Scientifique.



a1)  $a(x, -t, -\xi) = a(x, t, \xi)$  for a.a.  $x$  in  $\Omega$ , all  $(t, \xi)$  in  $\mathbb{R} \times \mathbb{R}^N$ .

a2)  $a(x, 0, 0) = 0$  for a.a.  $x$  in  $\Omega$ .

If we denote by  $a_i(x, t, \xi)$ ,  $i = 1, \dots, N$ , the partial derivative of  $a$  with respect to  $\xi_i$  and  $a_0(x, t, \xi)$  the partial derivative of  $a$  with respect to  $t$ , we assume:

a3) there exists  $C > 0$  and  $2 \leq p < \infty$  such that:

$$|a_\alpha(x, t, \xi)| \leq C(k(x) + |t|^{p-1} + |\xi|^{p-1}), \text{ where } |\xi| =$$

$$\left| \sum_{i=1}^N \xi_i^2 \right|^{\frac{1}{2}} \text{ for } \alpha = 0, 1, \dots, N, \text{ and } k \in L^{p/p-1}(\Omega).$$

a4) Leray-Lions conditions:

$$i) \sum_{i=1}^N [a_i(x, t, \xi) - a_i(x, t, \xi')] (\xi_i - \xi'_i) > 0 \text{ for } \xi \neq \xi'$$

$$ii) \lim_{|\xi| \rightarrow +\infty} \left( \sum_{i=1}^N a_i(x, t, \xi) \xi_i \right) / (|\xi| + |\xi|^{p-1}) \rightarrow \infty$$

for a.a.  $x$  in  $\Omega$  and  $|t|$  bounded.

Let  $f: \Omega \times \mathbb{R} \rightarrow \mathbb{R}$  be measurable for all  $t \in \mathbb{R}$  and continuous for almost all  $x$  in  $\Omega$ , satisfying:

$$f1) \operatorname{ess\,sup}_{x \in \Omega} \sup_{|s| \leq t} |f(x, s)| \leq K(t), \text{ for all } t \in \mathbb{R}$$

$$f2) f(x, t)t \geq 0 \text{ for almost all } x \text{ in } \Omega.$$

Remark. If  $f$  doesn't depend on  $x$ ,  $f1)$  is trivially satisfied and  $f2)$  is purely a sign condition. Observe that by our assumptions on  $a$ ,  $\varphi(u) := \int_{\Omega} a(x, u, Du) dx$  is  $C^1(W_0^{1,p}, \mathbb{R})$ .

We consider the following equation:

$$(3) \quad \int_{\Omega} \sum_{i=1}^N a_i(x, u, Du) D^i v \, dx + \int_{\Omega} f(x, u) v \, dx = \lambda \int_{\Omega} u v \, dx$$

$$\text{for all } v \in W_0^{1,p}(\Omega) \cap L^{\infty}(\Omega) \text{ and } \int_{\Omega} |u|^2 \, dx = R^2 > 0 .$$

By a positive (resp. negative) solution of (3), we mean a pair  $(\lambda, u) \in \mathbb{R} \times W_0^{1,p}(\Omega)$  such that  $u$  is  $\geq 0$  a.e. (resp  $\leq 0$  a.e.),  $f(u)$  and  $f(u)u$  are in  $L^1(\Omega)$ , and satisfy (3) and (4):

$$(4) \quad \int_{\Omega} \sum_{i=1}^N a_i(x, u, Du) D^i u \, dx + \int_{\Omega} f(x, u) u \, dx = \lambda R^2 .$$

Our main result is:

Theorem 1. If  $a$  and  $f$  satisfy  $a_1 - a_4$  and  $f_1 - f_2$ , and if  $\lim_{\|u\|_{W_0^{1,p}} \rightarrow \infty} \varphi(u) = +\infty$ ,

then for all  $R > 0$ , (3) possesses at least one positive and one negative solution.

## 2. Proof.

a) First observe that if  $u \in W_0^{1,p}(\Omega)$ , then  $u^+ = \sup(u, 0)$  belongs to  $W_0^{1,p}(\Omega)$  as well as  $|u| = \sup(u, -u)$ . Let  $\Omega^+ \subset \Omega$  be the support (as a distribution) of  $u^+$ .

We have  $\varphi(|u|) = \varphi(u)$ . Indeed  $\varphi(|u|) = \int_{\Omega} a(x, |u|, D|u|) \, dx = \int_{\Omega^+} a(x, u, Du) \, dx + \int_{\Omega - \Omega^+} a(x, -u, -Du) \, dx = \int_{\Omega^+} a(x, u, Du) \, dx + \int_{\Omega - \Omega^+} a(x, u, Du) \, dx = \varphi(u)$ , by  $a_1$ .

b) Since we are looking for positive solutions, without loss of generality we can assume that  $f$  is odd. Indeed we can replace  $f$  by  $\tilde{f}$  defined by  $\tilde{f}(x, t) = f(x, t)$  for  $t \geq 0$  and  $\tilde{f}(x, t) = -f(x, -t)$  for  $t < 0$ . The negative case is similar.

By a1), if  $f$  is odd, and  $(\lambda, u)$  is a positive solution, then  $(\lambda, -u)$  is a negative one, so we can restrict ourselves to the case of positive solutions, with  $f$  odd.

c) We shall first prove the theorem under the additional assumption that  $f$  is

bounded, let  $\psi(u) := \int_{\Omega} dx \int_0^{u(x)} f(x, t) dt$ . It is known that  $\psi \in C^1(W_0^{1,p}, \mathbb{R})$ .

Clearly  $\psi \geq 0$ . By the compact imbedding of  $W_0^{1,p}(\Omega)$  in  $L^2(\Omega)$  ( $p \geq 2$ ), if

$u_n \rightharpoonup u$  in  $W_0^{1,p}$ , then  $u_n \rightarrow u$  in  $L^2$  and therefore  $\psi(u_n) \rightarrow \psi(u)$ . We know

that  $\varphi \in C^1(W_0^{1,p}, \mathbb{R})$ . Moreover  $\varphi'(u)$ , the Frechet derivative of  $\varphi$  at  $u$  is bounded by

a3) and satisfies [see 5, p. 183]:  $u_n \rightharpoonup u$  in  $W_0^{1,p}$ ,  $\varphi'(u_n) \rightharpoonup v$  in  $(W_0^{1,p})'$  and

$\overline{\lim} \langle \varphi'(u_n), u_n - u \rangle \leq 0$  imply  $v = \varphi'(u)$  and  $\lim \langle \varphi'(u_n), u_n \rangle = \langle v, u \rangle^\dagger$ .

It easily follows that  $\varphi': W_0^{1,p} \rightarrow (W_0^{1,p})'$  is of type  $(P)^\dagger$ , [6] and therefore

$\varphi: W_0^{1,p} \rightarrow \mathbb{R}$  is sequentially weakly lower semicontinuous. Hence  $\varphi + \psi$  is also s.w.l.s.

For  $R > 0$ , let  $\alpha := \inf_{v \in S^R} \varphi(v) + \psi(v)$ , where  $S^R := \{u \in W_0^{1,p} \mid \int_{\Omega} |u|^2 dx = R^2\}$ .

Clearly  $S^R \neq \emptyset$ , so  $\alpha < \infty$ . Let  $u_n \in S^R$  such that  $\varphi(u_n) + \psi(u_n) \downarrow \alpha$ .

Since  $\psi \geq 0$  and by the assumption of the theorem,  $\|u_n\|_{W_0^{1,p}} \leq C$ , for some  $c > 0$ .

We have  $\varphi(|u_n|) + \psi(|u_n|) = \varphi(u_n) + \psi(u_n)$  and  $\| |u_n| \|_{W_0^{1,p}} = \|u_n\|_{W_0^{1,p}} \leq C$ .

By the compact imbedding of  $W_0^{1,p}$  into  $L^2$ , and the reflexivity of  $W_0^{1,p}$  there

exists  $u \in W_0^{1,p}$  with  $\int_{\Omega} |u|^2 dx = R^2$  and  $u \geq 0$  such that  $|u_n| \rightharpoonup u$  in  $W_0^{1,p}$

and  $|u_n| \rightarrow u$  in  $L^2$ . Therefore, by our previous remark,  $\varphi(u) + \psi(u) \leq \underline{\lim} \varphi(|u_n|)$

$+ \psi(|u_n|) = \alpha$ . But  $u \in S^R$ , so  $\varphi(u) + \psi(u) = \alpha$ . Hence the minimum of

$\dagger \langle \cdot, \cdot \rangle$  shall denote the duality between  $W_0^{1,p}$  and  $(W_0^{1,p})'$ .

$\ddagger u_n \rightharpoonup u$  in  $W_0^{1,p}$  implies  $\overline{\lim} \langle \varphi'(u_n), u_n - u \rangle \geq 0$ .



$\varphi + \psi$  on  $S^R$ , is achieved at  $u$ . Moreover  $\varphi + \psi \in C^1(W_0^{1,p}, \mathbb{R})$ ,  $u \rightarrow \frac{1}{2} \int_{\Omega} |u|^2 dx$  is  $C^1(W_0^{1,p}, \mathbb{R})$  and the Frechet derivative of the latter function is  $\neq 0$  since  $u \neq 0$ . Thus, by the well-known "Lyusternik principle", there exists  $\lambda \in \mathbb{R}$ , such that  $(\lambda, u)$  is a positive solution of (3).

d) For  $n \in \mathbb{N}$ , define  $f_n(x, t) = f(x, t)$  if  $|f(x, t)| \leq n$ ,  $f_n(x, t) = n$  if  $f(x, t) > n$  and  $f_n(x, t) = -n$  if  $f(x, t) < -n$ . Let  $\psi_n(u) := \int_{\Omega} dx \int_0^{u(x)} f_n(x, t) dt$ . Since  $f_n$  are bounded, we know that for each  $n \in \mathbb{N}$ , there exists  $(\lambda_n, u_n) \in \mathbb{R} \times W_0^{1,p}$ , a positive solution of (3), where  $f$  is replaced by  $f_n$ . We claim that

$\|u_n\|_{W_0^{1,p}} \leq C$  for some  $C > 0$ . Indeed as in c), we have:  $\varphi(u_n) + \psi_n(u_n) = \inf_{v \in S^R} \varphi(v) + \psi_n(v)$ . Let  $u_0 \in S^R \cap L^\infty(\Omega)$ . Then  $\varphi(u_n) + \psi_n(u_n) \leq \varphi(u_0) + \psi_n(u_0) \leq \varphi(u_0) + \int_{\Omega} dx \int_0^{u_0(x)} f(x, t) dt < \infty$ . By the coercivity of  $\varphi$  and the positivity of  $\psi_n$  we get  $\|u_n\|_{W_0^{1,p}} \leq C$ . By the reflexivity of  $W_0^{1,p}$  and the compact imbedding of  $W_0^{1,p}$  into  $L^2$  we can extract a subsequence, still denoted by  $u_n$ , such that  $u_n \rightharpoonup u$  in  $W_0^{1,p}$  and  $u_n \rightarrow u$  in  $L^2$ . So  $u \in S^R$  and  $u$  is positive a.e.

We will prove that  $\lambda_n$  is bounded. Indeed, assume that there is a subsequence  $\lambda_n \downarrow -\infty$ , then we get:

$$(5) \quad \frac{1}{\lambda_n} \langle \varphi'(u_n), u_n \rangle + \frac{1}{\lambda_n} \int_{\Omega} f_n(x, u_n) u_n dx = R^2.$$

But  $\varphi'(u_n)$  is bounded, since  $u_n$  is, so  $\frac{1}{\lambda_n} \langle \varphi'(u_n), u_n \rangle \rightarrow 0$ . But then  $\frac{1}{\lambda_n} \int_{\Omega} f_n(x, u_n) u_n dx \leq 0$  and  $R^2 > 0$  imply that for  $n$  big enough we get a contradiction. Next assume now that  $\lambda_n \uparrow \infty$ , for a subsequence. We have, for all  $v \in W_0^{1,p}$ :

$$(6) \quad \frac{1}{\lambda_n} \langle \varphi'(u_n), v \rangle + \frac{1}{\lambda_n} \int_{\Omega} f_n(x, u_n) v dx = \int_{\Omega} u_n v dx .$$

In particular for  $v = u_n$  we get:

$$(7) \quad \int_{\Omega} \frac{1}{\lambda_n} f_n(x, u_n) u_n dx \leq C \text{ for some } C > 0 .$$

By assumption f1) and f2), for each  $\delta > 0$ , there exists  $K_{\delta} > 0$  such that

$$|f(x, t)| \leq K_{\delta} + \delta f(x, t)t . [2]. \text{ Hence } |f_n(x, t)| \leq K_{\delta} + \delta f(x, t)t . \text{ By using (7),}$$

this shows that the sequence  $\frac{1}{\lambda_n} f_n(x, u_n)$  is equiintegrable and since

$\frac{1}{\lambda_n} f_n(x, u_n) \rightarrow 0$  a.e. in  $\Omega$ , by Vitali's theorem,  $\frac{1}{\lambda_n} f_n(x, u_n) \rightarrow 0$  in  $L^1$  and by (6), for  $v \in W_0^{1,p}(\Omega) \cap L^{\infty}(\Omega)$ , we get:  $\int_{\Omega} u v dx = 0$ . Hence by density,

$\int_{\Omega} |u|^2 dx = 0$ , a contradiction. Thus  $\lambda_n$  is bounded and we can extract a subsequence  $\lambda_n \rightarrow \lambda \in \mathbb{R}$ .

Since  $u_n$  is bounded in  $W_0^{1,p}$  and  $\varphi': W_0^{1,p} \rightarrow (W_0^{1,p})'$  is bounded,  $\varphi'(u_n)$  is bounded and we can extract a subsequence converging weakly to  $w \in (W_0^{1,p})'$ :  $\varphi'(u_n) \rightharpoonup w$  in  $(W_0^{1,p})'$ . We have  $\int_{\Omega} f_n(x, u_n) u_n dx = \lambda_n R^2 - \langle \varphi'(u_n), u_n \rangle \leq C$ . By using the same argument as before, there exists a subsequence  $f_n(x, u_n) \rightarrow f(x, u)$  in  $L^1$ . Moreover, since  $f_n(x, u_n) u_n \geq 0$ , by Fatou's lemma,  $f(x, u) u \in L^1$ . Therefore for all  $v \in W_0^{1,p} \cap L^{\infty}$ , we have:

$$(8) \quad \langle w, v \rangle + \int_{\Omega} f(x, u) v dx = \lambda \int_{\Omega} u v dx .$$

Now, we define  $v_n := \inf(u, n)$ . It is known that  $v_n \in W_0^{1,p} \cap L^{\infty}$ ,  $v_n \rightarrow u$  in  $W_0^{1,p}$  and by Lebesgue's dominated convergence theorem and the fact that  $f(x, u) u \in L^1$ ,  $f(x, u) v_n \rightarrow f(x, u) u$  in  $L^1$ . By putting  $v = v_n$  and passing to the limit we have:

$$(9) \quad \langle w, u \rangle + \int_{\Omega} f(x, u) u dx = \lambda R^2.$$

Now, we are done provided that we prove that  $w = \varphi'(u)$ . But  $u_n \rightarrow u$  in  $W_0^{1,p}$ ,  $\varphi'(u_n) \rightarrow w$  in  $(W_0^{1,p})'$ , so it is sufficient to prove that

$$\begin{aligned} \overline{\lim} \langle \varphi'(u_n), u_n - u \rangle &\leq 0. \text{ We have } \overline{\lim} \langle \varphi'(u_n), u_n - u \rangle = \overline{\lim} \langle \varphi'(u_n), u_n \rangle \\ &- \langle w, u \rangle = \overline{\lim} \left[ \lambda_n R^2 - \int_{\Omega} f_n(x, u_n) u_n dx \right] - \left[ \lambda R^2 - \int_{\Omega} f(x, u) u dx \right] = \int_{\Omega} f(x, u) u dx \\ &- \lim \int_{\Omega} f(x, u) u dx - \lim \int_{\Omega} f_n(x, u_n) u_n dx \leq 0, \text{ by Fatou's lemma. This completes} \\ &\text{the proof of the theorem.} \quad \square \end{aligned}$$

Remark 1. With the same arguments, we can allow a more general right hand side and consider the equation:

$$(10) \quad \int_{\Omega} \sum_{i=1}^N a_i(x, u, Du) D^i v dx + \int_{\Omega} f(x, u) v dx = \lambda \int_{\Omega} g(x, u) v dx$$

for all  $v \in W_0^{1,p} \cap L^{\infty}$  with the condition  $\int_{\Omega} |u|^2 dx = R^2 > 0$  replaced by the condition  $\int_{\Omega} dx \int_0^{u(x)} g(x, t) dt = R > 0$ . In this case if  $g: \Omega \times \mathbb{R} \rightarrow \mathbb{R}$  satisfies the Caratheodory condition, and

g1)  $g(x, t)t > 0$  for a. a.  $x$  in  $\Omega$

g2) there exists  $1 < q < \infty$  and  $C > 0$  such that  $|g(x, t)| \leq C(1 + |t|^{q-1})$

and if we assume that  $p$  in the hypothesis a3) satisfies  $1 < p < \infty$ , we can state the following generalization of Theorem 1:

Theorem 1'. If  $a, f, g$  satisfy  $a_1 - a_4$ ,  $f_1 - f_2$ ,  $g_1 - g_2$  and if we have

i) the injection of  $W^{1,p}$  into  $L^q$  is compact

ii)  $\lim_{\|u\|_{W^{1,p}} \rightarrow +\infty} \int_{\Omega} a(x, u, Du) dx = +\infty$



$$\text{iii)} \quad \lim_{\|u\|_{L^q} \rightarrow \infty} \int_{\Omega} dx \int_0^{u(x)} g(x, t) dt = +\infty.$$

Then the equation (10) possess at least one positive and one negative solution  $(\lambda, u) \in W_0^{1,p}(\Omega)$  with  $f(x, u)$  and  $f(x, u)u \in L^1$ , for all  $R > 0$ .

Remark 2. In the simple case  $-\Delta u + f(u) = \lambda u$ ,  $u \in W_0^{1,2}(\Omega)$ , our condition on  $f$  reduces to: i)  $f(0) = 0$  ii)  $\lim_{t \rightarrow +\infty} \frac{f(t)}{t} \geq -c$  for some  $c > 0$ . If we assume further that  $f(u) = \tilde{f}(u)u$ ,  $\tilde{f}(t) \geq 0$ ,  $\tilde{f} \in C^1(\mathbb{R})$ , it follows from [7], that there exists an unbounded connected set  $C^+$  of positive solutions in  $\mathbb{R} \times C^{1,\alpha}(\Omega)$  ( $0 < \alpha < 1$ ), containing  $(\lambda_0, 0)$  in its closure, where  $\lambda_0$  is the first eigenvalue of  $-\Delta h + \tilde{f}(0)h = \lambda h$ ,  $h|_{\partial\Omega} = 0$ . If we suppose that  $\lim_{t \rightarrow \infty} \tilde{f}(t) = +\infty$ , it follows easily that the projection of  $C^+$  on  $\mathbb{R}$  contains the interval  $[\lambda_0, \infty)$ . Therefore in this case the above equation possesses positive solutions with arbitrary norm in  $L^2(\Omega)$  and for all  $\lambda \geq \lambda_0$ . It can happen that all positive solutions are in  $C^+$ ; it is true, for example, if  $\tilde{f}'(t) \geq 0$  for all  $t \in \mathbb{R}$ . For the case  $\tilde{f}'(t) > 0$ , see [8]; the case  $\tilde{f}'(t) \geq 0$  can be handled by using a similar argument to [9]. In this particular situation  $C^+$  is a  $C^1$  curve of solutions in  $C^{2,\alpha}(\Omega)$ , parametrized by  $\lambda$  and unbounded in  $L^2(\Omega)$ . Concerning the regularity of the solutions, let us mention that if  $f \in C^0(\mathbb{R})$ ,  $f(0) = 0$  and  $f$  is monotone, then  $u \in C^{1,\alpha}(\Omega)$ . For a different approach of this case, see [10].



# REFERENCES

- [1] F. E. Browder, Existence theory for boundary value problems for quasi-linear elliptic systems with strongly nonlinear lower order terms (Proc. Sympos. Pure Math., vol. 23, 1973, p. 269-286).
- [2] P. Hess, Variational inequalities for strongly nonlinear elliptic operators (J. Math. Pures et Appl., t. 52, 1973, p. 285-298).
- [3] P. Hess, A strongly nonlinear elliptic boundary value problem (J. Math. Anal. and Appl., vol. 43, 1973, p. 241-249).
- [4] N. P. Các, On strongly nonlinear variational inequalities (J. Math. Anal. and Appl., vol. 54, 1975, p. 1-10).
- [5] J.-L. Lions, Quelques méthodes de résolution des problèmes aux limites nonlinéaires, Dunod, Gauthier-Villars, Paris, 1969.
- [6] P. Hess, A variational approach to a class of nonlinear eigenvalue problems (Proc. Amer. Math. Soc., vol. 29, 1971, p. 272-276).
- [7] P. H. Rabinowitz, Some aspects of nonlinear eigenvalue problems (Rocky Mtn. J. of Math., 3, 1973, p. 161-202).
- [8] P. H. Rabinowitz, A note on a nonlinear eigenvalue problem for a class of differential equations (J. Diff. Eq., 9, 1971, p. 536-548).
- [9] Ph. Clément, Continuous branches of positive solutions for a class of nonlinear second-order differential equations (Manuscripta Math., 16, 1975, p. 195-202).

- [10] Ph. Clément, Nonlinear eigenvalue problem for a class of cyclically maximal monotone operators, to appear.



UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 1673	2. GOVT ACCESSION NO.	3. TECHNICAL DATA NUMBER (9) Technical
4. TITLE (and Subtitle) POSITIVE EIGENFUNCTIONS FOR A CLASS OF SECOND-ORDER ELLIPTIC EQUATIONS WITH STRONG NONLINEARITY.		5. TYPE OF REPORT & PERIOD COVERED Summary Report, no specific reporting period
7. AUTHOR(s) (10) Philippe Clement		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of 610 Walnut Street Wisconsin Madison, Wisconsin 53706		8. CONTRACT OR GRANT NUMBER(s) (15) DAAG29-75-C-0024
11. CONTROLLING OFFICE NAME AND ADDRESS U. S. Army Research Office P.O. Box 12211 Research Triangle Park, North Carolina 27709		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) (14) MRC-TSR-1673		12. REPORT DATE (11) Sep 1976
		13. NUMBER OF PAGES 10 (12) 13 p.
		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)  Nonlinear eigenvalue problem, Strong nonlinearity, Positive solution.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Let $\Omega$ be a bounded domain of $\mathbb{R}^N$ . We consider the equation $\mathcal{A}u(x) + f(x, u(x)) = \lambda u(x)$ , $\int_{\Omega}  u ^2(x) dx = R^2 > 0$ , where $\mathcal{A}$ is a second-order quasi-linear elliptic operator whose coefficients have polynomial growth and $f$ essentially satisfies a sign condition. The existence of positive and negative solutions is proved.		

DD FORM 1473  
1 JAN 73

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

22/200